

LIS00PP22245P3

(12) United States Plant Patent

(10) **Patent No.:**

US PP22,245 P3

(45) **Date of Patent:**

Nov. 15, 2011

(54) CHERRY TREE NAMED 'KORVIK'

(50) Latin Name: *Pruntus avium*Varietal Denomination: **Korvik**

(76) Inventor: Vaclav Ludvik, Horice (CZ)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 12/460,904

(22) Filed: Jul. 23, 2009

(65) **Prior Publication Data**

US 2010/0043110 P1 Feb. 18, 2010

Related U.S. Application Data

(60) Provisional application No. 61/137,623, filed on Jul. 31, 2008.

(51) **Int. Cl.** *A01H 5/00* (2006.01)

(52) U.S. Cl. Plt./181

(56) References Cited

OTHER PUBLICATIONS

GTITM UPOVROM Citation for 'Korvik' as per CZ PBR application filing No. 1327; Dec. 20, 2001.*

GTITM UPOVROM Citation for 'Korvik' as per CZ NLI application filing No. 6558; Dec. 20, 2001.*

* cited by examiner

Primary Examiner — Kent L Bell (74) Attorney, Agent, or Firm — Ballew Law

(57) ABSTRACT

'Korvik' is a new and distinct cherry variety (*Prunus avium*) notable for its round shape and resistance to cracking. 'Korvik' is further notable for its productivity on dwarfing rootstocks, and tolerance to Monilia disease ("brown rot").

2 Drawing Sheets

2

1

Latin name of the genus and species of the plant claimed: *Prumus avium*.

Variety denomination: 'Korvik'.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

None.

BACKGROUND AND SUMMARY OF THE VARIETY

'Korvik' originated in 1973 as a result of a controlled cross of 'Kordia' (female parent) and 'Vic' (male parent) cherry trees. Asexual propagation by budding was first carried out in 1975 in East Bohemia, Czech Republic. The variety has since been observed over a number of asexually propagated generations, and has been found to remain true to type.

'Korvik' is a new and distinct cherry variety notable for its round shape and resistance to cracking. 'Korvik' is further notable for its productivity on dwarfing rootstocks, and tolerance to Monilia disease ('brown rot'). 'Korvik' is more upright than its parent 'Kordia' which is more spreading and has less number of lenticels on the tree stem than its parent 'Vic'. The duration of the flowering of 'Korvic' is 1-2 days longer than its parent 'Vic'. In addition, 'Korvik' differs from parental varieties by leaf shape. The leaf margin of 'Korvik' is obtuse serrate, whereas the leaf margin of its parent, 'Kordia', is acute serrate.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

FIG. 1 shows the fruit and leaves of 'Korvik'; and

FIG. 2 shows the fruit of 'Korvik'.

DETAILED BOTANICAL DESCRIPTION

The following detailed botanical description is based on observations of 'Korvik' cherry trees grown at East Bohemia, ³⁵

Czech Republic. Color descriptions refer to The Royal Horticultural Society Colour Chart. It should be understood that

the characteristics described will vary somewhat depending upon cultural practices and climatic conditions, and can vary with location and season. Quantified measurements are expressed as an average of measurements taken from a number of individual plants of the new variety. The measurements of any individual tree, or any group of trees, of the new variety may vary from the stated average.

Tree:

Size.—Height: 3.9 m; Diameter 2.2 m.

Form.—Conical.

Vigor.—Medium.

Hardiness.—High.

Trunk.—Diameter: 96 mm at 0.5 m above ground; Texture: Medium smooth; Lenticels: very few in number; Color: greyed orange 165B.

Branches.—Scaffold branch length: 0.8 m; Diameter: 24 mm; Texture: smooth; Color: brown 200B; Number of lenticels per cm²: 4-5; Lenticel length: 2-3 mm; Lenticel diameter: 1 mm; Lenticel color greyed orange 165B; Fruiting branch length: 0.6 m; Diameter: 8.5 mm; Texture: smooth; Color: brown 200B; Number of lenticels per cm²: 7; Lenticel length: 1 mm; Lenticel diameter: 0.5 mm; Lenticel color: greyed orange 165B; Current season branch length: 0.6 m; Texture: smooth; Color: greyed orange 177B; Number of lenticels per cm²: 6; Lenticel length: 0.3 mm; Lenticel diameter 0.3 mm; Lenticel color: greyed orange 175B.

Leaf:

30

Size.—Length: 105 mm; Width: 62 mm. Shape.—Oval; Apex: pointed; Base: obtuse.

Margin.—Obtuse serrate.

3

Texture.—Upper surface: glossy; Lower surface: dull with marked veining.		Pollen amount.—Medium. Pollen color.—Yellow orange 14A.
Color.—Upper surface: green 136B; Lower surface: green 137C.		Carpel style length.—12 mm. Carpel style color.—Yellow green 145D.
Petiole.—Length: 22 mm; Diameter: 0.4 mm; Color: greyed purple 183B.	5	Style color.—Yellow green 145C. Stigma shape.—Oval.
Leaf glands:		Stigma diameter.—0.6 mm.
Size.—Length: 1 mm; Width: 0 8 mm.		Stigma color.—Yellow green 145B.
Shape.—Oval.		Fruit:
Number per leaf.—1-2.	10	Size.—Large; Height: 27.2 mm; Suture diameter: 28.3
Location.—Mostly on the petiole, upper position.		mm; Apex diameter: 22.7 mm; Stem cavity: Medium,
Color.—Greyed purple 183A.		Depth: 2-2.5 mm, Breadth 10-11 mm; Peduncle:
Flower buds:		Length 48-55 mm, diameter 1.3-1.5 mm, Color:
Size.—Length: 6.5 mm; Diameter: 3.2 mm.		Green 146C; Quantity of fruit per cluster: 2-4.
Shape.—Conical.	15	Fruit shape.—Round; Base shape: Round to oval.
Number on each fruiting spur.—2.5.		Skin.—Color: Greyed purple 187B; Thickness:
Color.—Greyed purple 166A.		Medium, limited tendency to cracking.
Blooming period.—14 days.		Flesh.—Color Greyed purple 187D; Flesh Texture:
Date of full bloom.—Medium late (April 22).		Above mean 6 (if the rating scale from 1 worst to 9
Flowers:	20	best is used); Flesh Fiber: Medium; Flavor: Medium,
Petal size.—Length: 15 mm; Width: 12 mm.		slightly acidulated sweet; Aroma: Good; Eating qual-
		ity: Good; Brix: Varies from 16.2° to 20.6°.
Petal number.—5 per flower.		Stone.—Size: Length 12.5 mm, Width 10.5 mm at its
Petal shape.—Oval; Apex: rounded; Base: slightly		widest point near the center; Form: Oblong; Sides:
pointed.	25	Ridges 4-6 with varying amounts of encircling the
Petal margin.—Smooth.		sutures of the pit; No tendency to pit cracking; Color
Petal texture.—Upper surface: slight venous; Lower surface: nearly smooth.		Orange white 159A; Kernal size; Length 8.9 mm, Width at widest point: 5.6 mm; Kernel color: Orange
Petal color.—Upper surface: white N155A; Lower sur-		22A.
face: red-purple 69C.	30	Firmness.—Medium; It was measured by Durofel DTT
Sepals:		100 Agrotechnologie SA-Licence CTIFL (Digital
Size.—Length: 6 mm; Diameter: 4 mm.		Firmness Tester France); Its scale values ranged from
Number.—5.		$0-99$ where $85=4.5 \text{ kg/}0.5 \text{ cm}^2$ of penetrometer mea-
Shape.—Widely conical; Apex: pointed.		surement; Mean value of 'Korvik' was 69.
Margin.—Smooth.	35	Fruit weight.—2nd Year after planting: 0.25 kg; 3 rd Year
Texture.—Upper surface: dull smooth; Lower surface:		after planting 1 kg; 4 th Year after planning: 5 kg.
glossy smooth.		Time of ripening.—Late.
Color.—Upper surface: Yellow green 147B; Lower sur-		Harvest.—Started on July 11 th in 2010 in East Bohemia,
face: green 138B.		Czech Republic.
Peduncle length.—36 mm; Diameter: 1 mm; Color:	40	Orchard performance:
green 137B.		Disease resistance/tolerance.—Tolerant to Monilia and
Reproductive characteristics:		Blumeriella jaapii infection.
Nectary color.—Yellow 2C.		Yield.—Very good yields on dwarf rootstocks.
Stamen number.—16.		I claim:
Filament color.—Red-purple 69C.	45	1. A new and distinct cherry tree substantially as shown and
Anther shape.—Round to oval.		described herein.
Anther size.—Length: 0.5 mm; Diameter: 0.5 mm.		
Anther color.—Yellow orange 14A.		* * * *



FIG. 1




FIG. 2